

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/09668 A2

- (51) International Patent Classification⁷: **A61K 9/00** SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/US01/21105
- (22) International Filing Date: 20 July 2001 (20.07.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/222,476 2 August 2000 (02.08.2000) US
- (71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MICANOVIC, Radmila** [US/US]; 7126 White Oak Trail, Indianapolis, IN 46236 (US). **WITCHER, Derrick, Ryan** [US/US]; 10898 Parrot Court, Fishers, IN 46038 (US).
- (74) Agents: **WEBSTER, Thomas, D.** et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- Published:**
- without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/09668 A2

(54) Title: PULMONARY ADMINISTRATION OF FLINT

(57) Abstract: The claimed invention relates to methods and compositions for administering FLINT by inhalation, and a methods for treating a disease by administering FLINT by inhalation.

-1-

PULMONARY ADMINISTRATION OF FLINT**Field of the Invention**

This invention relates to the pulmonary delivery of
5 FLINT for systemic absorption through the lungs to
significantly reduce or eliminate the need for administering
FLINT by injection or other routes of administration.

Background of the Invention

10 A number of proteins homologous to tumor necrosis
factor receptor proteins ("TNFR proteins") have been
isolated in recent years, having many potent biological
effects. Aberrant activity of these proteins has been
implicated in a number of disease states.

15 A recently discovered TNFR homologue, referred to
herein as "FAS Ligand Inhibitory Protein," or "FLINT", binds
Fas Ligand (FasL) to prevent the interaction of FasL with
Fas (See PCT Application No. WO 99/50413).

Increased activation of the Fas-FasL signal
20 transduction pathway is thought to underlie a number of
pathological conditions, including runaway apoptosis (Kondo
et al., Nature Medicine 3(4):409-413 (1997); Galle et al.,
J. Exp. Med. 182:1223-1230 (1995)), and inflammatory disease
resulting from neutrophil activation (Miwa et al., Nature
25 Medicine 4:1287 (1998)).

Runaway apoptosis signifies a level of apoptosis
greater than normal, or apoptosis occurring at an
inappropriate time. Pathological conditions caused by
runaway apoptosis include, for example, organ failure in the
30 liver, kidneys and pancreas. Inflammatory diseases
associated with excessive neutrophil activation include
sepsis, ARDS, SIRS and MODS.

-2-

FLINT inhibits not only the binding of Fas to FasL, but also the binding of another ligand, LIGHT, to LT β R, and/or TR2/HVEM receptors. This property can be used to treat or prevent diseases or conditions that may be associated with these binding interactions.

Like many other proteins, FLINT is thought to rapidly degrade in the gastrointestinal tract, and therefore FLINT is not efficiently absorbed from the GI tract into the blood stream. Therefore, exogenous administration of FLINT for therapeutic purposes depends, in most instances, upon injection treatments. Alternatively, some proteins may be efficiently administered by rectal, transdermal, and nasal routes. Thus far, however, these routes of administration have not resulted in generally effective protein absorption.

Over the years, certain drugs have been sold in compositions suitable for forming a drug dispersion for oral inhalation (pulmonary delivery) to treat various conditions in humans. Such pulmonary drug delivery compositions are designed to be delivered by inhalation of a drug dispersion by the patient so that the active drug within the dispersion can reach the lung. It has been found that certain drugs delivered to the lung are readily absorbed through the alveolar region directly into blood circulation.

Pulmonary delivery is particularly promising for the delivery of proteins and polypeptides which are difficult to deliver by other routes of administration. Such pulmonary delivery can be effective both for systemic delivery and for localized delivery to treat diseases of the lungs.

Pulmonary drug delivery can be achieved by different approaches, including liquid nebulizers, aerosol-based metered dose inhalers (MDI's), and dry powder dispersion devices. Dry powder dispersion devices, which do not rely on

CFC aerosol technology, are promising for delivering drugs that may be readily formulated as dry powders, particularly proteins and polypeptides. Many otherwise labile proteins and polypeptides may be stably stored as lyophilized or
5 spray-dried powders by themselves or in combination with suitable powder carriers.

The ability to deliver proteins and polypeptides as dry powders, however, is problematic in certain respects. The dosage of many protein and polypeptide drugs is often
10 critical so it is necessary that any dry powder delivery system be able to accurately, precisely, and reliably deliver the intended amount of drug. Moreover, many proteins and polypeptides are quite expensive, typically being many times more costly than conventional drugs on a per-dose
15 basis. Thus, the ability to efficiently deliver the dry powders with a minimal loss of drug is critical.

It is now known that effective absorption of proteins can occur from the lung. For example, administration of insulin as an inhalation aerosol to the lung was first
20 reported by Gaensslen in 1925. Recently, pulmonary administration of erythropoietin has been described (United States Patent No. 5,354,934).

The efficiency of pulmonary administration depends on numerous factors. To a large extent, pulmonary absorption is
25 dependent on the physical characteristics of the particular therapeutic protein to be delivered. Moreover, efficacy is at least partly dependent on the ability to deliver the protein to the deep lung alveolar epithelium. Proteins that are deposited in the upper airway epithelium are not
30 absorbed to a significant extent. This is due to the overlying mucus which is approximately 30 μm - 40 μm thick, which acts as a barrier to absorption. In addition,

-4-

proteins deposited on this epithelium are cleared by mucociliary transport up the airways and then eliminated via the gastrointestinal tract.

FLINT, as a therapeutic protein, is useful in treating
5 and/or preventing diseases that may relate to enhanced apoptosis. Therapeutic administration of FLINT can be accomplished by any number of suitable methods, including, for example, intravenous and subcutaneous administration. However, such treatments can be painful to a patient,
10 especially to one who must receive multiple dosings for effective treatment. Thus, there is a need for a route of therapeutic administration of FLINT that minimizes the pain and discomfort of injection methods.

15 **Summary of the Invention**

The present invention solves this problem by administering FLINT comprising administering by pulmonary means an effective amount of FLINT to a patient in need thereof.

20 The present invention also relates to a method for treating a disease related to enhanced apoptosis comprising administering by pulmonary means an effective dose of FLINT to a patient in need thereof. Preferably, FLINT is delivered by inhalation, to the lower airway of the patient.
25 A further object of this invention is to provide a powdered pharmaceutical composition containing FLINT suitable for pulmonary delivery.

The FLINT compounds of the invention can be delivered to the lung in a carrier, as a solution or suspension, or as
30 a dry powder, using any of a variety of devices suitable for pulmonary administration by inhalation. Preferably, a FLINT compound is delivered at a particle size effective for

-5-

reaching the lower airways of the lung. A preferred particle size is below 10 microns. An even more preferred particle size is between 1 and 5 microns.

5

Detailed Description of the Invention

The term "powder" or "powdered" refers to a composition that consists of finely dispersed solid particles that are relatively free flowing and capable of being dispersed in an inhalation device and subsequently inhaled by a subject so that the particles reach the lungs to permit penetration into the alveoli. Thus, the powder is administrable by inhalation therapy and is said to be "respirable" and suitable for pulmonary delivery. In general, the average particle size is less than about 10 microns (μm) in diameter and the particle shapes may be irregular, uniform or mixed. Preferably, the average particle size is less than about 7.5 μm and more preferably less than about 5.0 μm . Usually the particle size distribution is between about 0.1 μm and about 5 μm in diameter, particularly between about 2 μm to about 5 μm .

The term "dry" means that the powder composition has a moisture content such that the particles are readily dispersible in an inhalation device to form an aerosol. This moisture content is generally below about 10% by weight (%w) water, usually below about 5%w and preferably less than about 3%w.

The term "dispersibility" means the degree to which a powder composition can be dispersed (i.e. suspended) in a current of air so that the dispersed particles can be respired or inhaled into the lungs of a subject. For example, a powder composition that is only 10% dispersible means that only 10% of the mass of finely-divided particles

-6-

making up the composition can be suspended for inhalation into the lungs; 50% dispersibility means that 50% of the mass can be suspended. A standard measurement of dispersibility is described hereinafter.

5 The term "therapeutically effective amount" is the amount of FLINT present in the powder composition that is needed to provide the desired level to a subject to be treated to give the anticipated physiological response.

10 The term "physiologically effective amount" is that amount delivered to a subject to give the desired palliative or curative effect. This amount is specific for each active agent and its ultimately approved dosage level.

15 The term "pharmaceutically acceptable" refers to an excipient, whether a carrier or a protein used to improve dispersibility, that can be taken into the lungs with no significant adverse toxicological effects on the lungs.

20 The term "FLINT" as used herein refers to native FLINT (SEQ ID NO:3), mature FLINT lacking the leader sequence (SEQ ID NO:1) and/or to FLINT analogs. The term "protease-resistant analog" or "protease-resistant FLINT analog" means a variant FLINT sequence, or fragment thereof, resistant to proteolysis between positions 218 and 219 of SEQ ID NO:1 (positions 247 and 248 of SEQ ID NO:3). Protease resistant analogs may have one, two, three, or more amino acid
25 substitutions, deletions, additions, or other changes to the sequence of SEQ ID NO:1 or SEQ ID NO:3. Preferably, there is a single substitution in said sequence.

30 The term "FLINT analog" or "analog" refers to a variant FLINT, or fragment thereof, preferably having a biological activity substantially the same as FLINT, for example, the ability to inhibit apoptosis in vitro and/or in vivo. A FLINT analog comprises one or more substitutions, additions,

-7-

deletions in the amino acid sequence embodied in SEQ ID NO:1 or 3.

The term "protease-resistant" refers to a FLINT analog that, when compared with FLINT, or FLINT fragment, exhibits
5 resistance to proteolysis between residues 218 and 219 of SEQ ID NO:1. Protease resistant analogs differ from FLINT by one or more amino acid substitutions, deletions, inversions, additions, and/or changes in glycosylation sites, or patterns, as compared with native FLINT, or mature FLINT, or
10 FLINT fragment. Preferably these changes occur in the region from about position 214 through position 222 of SEQ ID NO:1.

The term "protease-resistant" contemplates degrees of resistance to proteolysis at position 218 from complete resistance to partial resistance. Thus, a "resistant analog"
15 may show a degree of resistance to proteolysis at position 218, as measured by half-life, that is at least about 25% greater than native FLINT when treated or exposed to a suitable protease. Preferably, a resistant FLINT analog possesses a protease resistance half-life that is at least
20 about 2-fold greater than native FLINT.

As used herein "half-life" refers to the time required for approximately half of FLINT or a FLINT analog molecule in a sample to be proteolytically cleaved between positions 218 and 219 of SEQ ID NO:1, *in vitro* and/or *in vivo*, as
25 determined by any suitable means.

Exemplary protease-resistant FLINT analogs include the following substitutions in SEQ ID NO:1:

- a. Arg at position 218 is replaced by Gln;
- b. Arg at position 218 is replaced by Glu;
- 30 c. Thr at position 216 is replaced by Pro;
- d. Arg at position 218 is replaced by Ala;
- e. Arg at position 218 is replaced by Gly;

-8-

- f. Arg at position 218 is replaced by Ser;
- g. Arg at position 218 is replaced by Val
- h. Arg at position 218 is replaced by Tyr;
- i. Pro at position 217 is replaced by Tyr
- 5 j. Thr at position 216 is replaced by Pro, and Arg at position 218 is replaced by Gln.

Additional protease-resistant FLINT analogs are disclosed in PCT Patent Application Serial No. PCT/US 00/06418.

The term "preservative" refers to a compound added to a pharmaceutical formulation to act as an anti-microbial agent. A parenteral formulation must meet guidelines for preservative effectiveness to be a commercially viable multi-use product. Among preservatives known in the art as being effective and acceptable in parenteral formulations are benzalkonium chloride, benzethonium, chlorohexidine, phenol, m-cresol, benzyl alcohol, methylparaben, chlorobutanol, o-cresol, p-cresol, chlorocresol, phenylmercuric nitrate, thimerosal, benzoic acid, and various mixtures thereof. See, e.g., Wallhäusser, K.-H., *Develop. Biol. Standard*, 24: 9-28 (Basel, S. Krager, 1974).

The term "buffer" or "pharmaceutically acceptable buffer" refers to a compound that is safe for use in FLINT formulations and has the effect of controlling the pH of the formulation at the pH desired for the formulation. Pharmaceutically acceptable buffers for controlling pH at a moderately acid pH to a moderately basic pH include, for example, such compounds as phosphate, acetate, citrate, TRIS, arginine, or histidine.

The term "isotonicity agent" refers to a compound that is tolerated physiologically and imparts a suitable tonicity to a formulation to prevent the net flow of water across the cell membrane. Compounds such as glycerin are commonly used

-9-

for such purposes at known concentrations. Other acceptable isotonicity agents include physiological solutions of salts, e.g., NaCl, dextrose, mannitol, and lactose. Glycerol at a concentration of 12 to 25 mg/mL is preferred as an

5 isotonicity agent.

Pulmonary Administration of FLINT

FLINT is administered by inhalation in a dose effective manner to increase circulating FLINT protein levels.

10 Administration of FLINT can be effective for treating a variety of disorders such as, for example, inflammatory diseases associated with neutrophil activation including sepsis, acute respiratory distress syndrome, acute lung injury, systemic inflammatory response syndrome (SIRS) and

15 multiple organ dysfunction (MODS).

Other diseases for which FLINT could be therapeutically effective include rheumatoid arthritis (Elliott et al., *Lancet* 344:1105-10 (1994)), fibroproliferative lung disease, fibrotic lung disease, HIV (Dockrell et al., *J. Clin. Invest.* 101:2394-2405 (1998)), ischemia (Sakurai et al. 1998 *Brain Res* 797:23-28), brain trauma/injury (Ertel et al. 1997 *J Neuroimmunol* 80:93-6), chronic renal failure (Schelling et al. 1998 *Lab Invest* 78:813-824), Graft-vs-Host Disease (GVHD) (Hattori et al. 1998 *Blood* 11:4051-4055), Cutaneous inflammation (Orteu et al. 1998 *J Immunol* 161:1619-1629), Vascular leak syndrome (Rafi et al. 1998 *J Immunol* 161:3077-3086), *Helicobacter pylori* infection (Rudi et al. 1998 *J Clin Invest* 102:1506-1514), Goiter (Tamura et al. 1998 *Endocrinology* 139:3646-3653), Atherosclerosis (Sata and Walsh, 1998 *J Clin Invest* 102:1682-1689), IDDM (Itoh et al. 1997 *J Exp Med* 186:613-618), Osteoporosis (Jilka et al. 1998 *J Bone Min Res* 13:793-802), Crohn's

20

25

30

-10-

Disease (van Dullemen et al. 1995 Gastroenterology 109:129-35), organ preservation and transplant (graft) rejection (Lau et al. 1996 Science 273:109-112), Sepsis (Faist and Kim. 1998 New Horizons 6:S97-102), Pancreatitis
5 (Neoptolemos et al. 1998 Gut 42:886-91), Cancer (melanoma, colon and esophageal) (Bennett et al. 1998 J Immunol 160:5669-5675), Autoimmune disease (IBD, psoriasis, Down's Syndrome (Seidi et al., Neuroscience Lett. 260:9 (1999) and multiple sclerosis (D'Souza et al. 1996 J Exp
10 Med 184:2361-70); Alzheimer's Disease; End-stage renal disease (ESRD); mononucleosis; EBV; Herpes; antibody dependent cytotoxicity; hemolytic and hypercoagulation disorders such as vascular bleeds, DIC (disseminated intravascular coagulation), eclampsia, HELLP (preeclampsia
15 complicated by thrombocytopenia, hemolysis and disturbed liver function), HITS (heparin induced thrombocytopenia), HUS (hemolytic uremic syndrome), and preeclampsia; hematopoietic disorders such as aplastic anemia, thrombocytopenia (TTP) and myelodysplasia; and hemolytic
20 fever caused, for example, by E.bola.

Achieving effective doses of FLINT requires administration of an inhaled dose of more than about 0.5 µg/kg to about 500 µg/kg FLINT protein; preferably about 3 µg/kg to about 250 µg/kg; and more preferably about 5 µg/kg
25 to about 100 µg/kg; more preferably still from about 7 µg/kg to about 50 µg/kg; still more preferably from about 7 µg/kg to about 20 µg/kg; most preferably from about 7 µg/kg to about 14 µg/kg. A therapeutically effective amount of FLINT can be determined by a knowledgeable medical practitioner
30 taking into account factors including the physical condition of the patient, the patient's pulmonary status, and the like.

-11-

According to the invention, FLINT is delivered by inhalation to achieve absorption of the protein. Administration by inhalation can result in pharmacokinetics comparable to subcutaneous administration. Different
5 inhalation devices typically provide similar pharmacokinetics when similar particle sizes and similar levels of lung deposition are compared.

According to the invention, FLINT can be delivered by any of a variety of inhalation devices known in the art for
10 administration of a therapeutic agent by inhalation. These devices include metered dose inhalers, nebulizers, dry powder generators, sprayers, and the like. Preferably, FLINT is delivered by a dry-powder inhaler, or sprayer. There are several desirable features of an inhalation device
15 for administering FLINT. For example, delivery by the inhalation device is advantageously reliable, reproducible, and accurate. The inhalation device should deliver small particles, e.g. less than about 10 μm , preferably about 1-5 μm , for good respirability. Some specific examples of
20 commercially available inhalation devices suitable for the practice of this invention are TurbohalerTM (Astra), Rotahaler[®] (Glaxo), Diskus[®] (Glaxo), SpirosTM inhaler (Dura), devices marketed by Inhale Therapeutics, AERxTM (Aradigm), the Ultravent[®] nebulizer (Mallinckrodt), the Acorn II[®]
25 nebulizer (Marquest Medical Products), the Ventolin[®] metered dose inhaler (Glaxo), the Spinhaler[®] powder inhaler (Fisons), or the like.

As those skilled in the art will recognize, the formulation of FLINT protein, the quantity of the
30 formulation delivered, and the duration of administration of a single dose depend on the type of inhalation device employed. For some aerosol delivery systems, such as

-12-

nebulizers, the frequency of administration and length of time for which the system is activated will depend mainly on the concentration of FLINT in the aerosol. For example, shorter periods of administration can be used at higher
5 concentrations of protein in the nebulizer solution.

Devices such as metered dose inhalers can produce higher aerosol concentrations, and can be operated for shorter periods to deliver the desired amount of protein. Devices such as powder inhalers deliver active agent until a given
10 charge of agent is expelled from the device. In this type of inhaler, the amount of FLINT protein in a given quantity of the powder determines the dose delivered in a single administration.

The particle size of the FLINT protein in the
15 formulation delivered by the inhalation device is important with respect to the ability of protein to make it into the lungs, and preferably into the lower airways or alveoli. Preferably, the FLINT is formulated so that at least about 10% of the protein delivered is deposited in the lung,
20 preferably about 10% to about 20%, or more. It is known that the maximum efficiency of pulmonary deposition is obtained with particle sizes of about 2 μm to about 3 μm . When particle sizes are above about 5 μm , pulmonary deposition decreases substantially. Particle sizes below
25 about 1 μm cause pulmonary deposition to decrease, and it becomes difficult to deliver particles with sufficient mass to be therapeutically effective. Thus, particles of FLINT protein delivered by inhalation have a particle size preferably less than about 10 μm , more preferably in the
30 range of about 1 μm to about 5 μm , and most preferably in the range of about 2 μm to about 3 μm . The formulation of

-13-

FLINT protein is selected to yield the desired particle size in the chosen inhalation device.

Administration of FLINT by a Dry Powder Inhaler

5 Advantageously for administration as a dry powder, FLINT is prepared in a particulate form with a particle size of less than about 10 μm , preferably about 1 to about 5 μm , and most preferably about 2 μm to about 3 μm . The preferred particle size is effective for delivery to the alveoli of
10 the patient's lung. Preferably, the dry powder is largely composed of particles produced so that a majority of the particles have a size in the desired range. Advantageously, at least about 50% of the dry powder is made of particles
15 having a diameter less than about 10 μm . Such formulations can be achieved by spray drying, milling, or critical point condensation of a solution containing FLINT and other desired ingredients. Other methods also suitable for generating particles useful in the current invention are known in the art.

20 The particles are usually separated from a dry powder formulation in a container and then transported into the lung of a patient via a carrier air stream. Typically, in current dry powder inhalers, the force for breaking up the solid is provided solely by the patient's inhalation. One
25 suitable dry powder inhaler is the Turbohaler[™] manufactured by Astra (Södertälje, Sweden). In another type of inhaler, air flow generated by the patient's inhalation activates an impeller motor which deagglomerates the protein particles. The Dura Spiros[™] inhaler is such a device.

30 Formulations of FLINT for administration from a dry powder inhaler typically include a finely divided dry powder containing FLINT protein, but the powder can also include a

-14-

bulking agent, carrier, excipient, another additive, or the like. Additives can be included in a dry powder formulation of protein, for example, to dilute the powder as required for delivery from the particular powder inhaler, to

5 facilitate processing of the formulation, to provide advantageous powder properties to the formulation, to facilitate dispersion of the powder from the inhalation device, to stabilize the formulation (e.g., antioxidants or buffers), to provide taste to the formulation, or the like.

10 Advantageously, the additive does not adversely affect the patient's airways. The FLINT protein can be mixed with an additive at a molecular level or the solid formulation can include particles of the FLINT protein mixed with or coated on particles of the additive. Typical additives include

15 mono-, di-, and polysaccharides; sugar alcohols and other polyols, such as, for example, lactose, glucose, raffinose, melezitose, lactitol, maltitol, trehalose, sucrose, mannitol, starch, or combinations thereof; surfactants, such as sorbitols, diphosphatidyl choline, or lecithin; or the

20 like. Typically an additive, such as a bulking agent, is present in an amount effective for a purpose described above, often at about 50% to about 90% by weight of the formulation. Additional agents known in the art for formulation of a protein can also be included in the

25 formulation.

Administration of FLINT as a Spray

A spray including FLINT protein can be produced by forcing a suspension or solution of FLINT protein through a

30 nozzle under pressure. The nozzle size and configuration, the applied pressure, and the liquid feed rate can be chosen to achieve the desired output and particle size. An

-15-

electrospray can be produced, for example, by an electric field in connection with a capillary or nozzle feed.

Advantageously, particles of FLINT delivered by a sprayer have a particle size less than about 10 μm , preferably in
5 the range of about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm .

Formulations of FLINT protein suitable for use with a sprayer typically include protein in an aqueous solution at a concentration of about 1 mg to about 20 mg of FLINT
10 protein per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the protein, such as a buffer, a reducing
15 agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating FLINT include albumin, protamine, or the like. Typical carbohydrates useful in formulating FLINT proteins include sucrose, mannitol, lactose, trehalose, glucose, or the like. The FLINT protein formulation can
20 also include a surfactant, which can reduce or prevent surface-induced aggregation of FLINT caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitol fatty
25 acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Additional agents known in the art for formulation of a protein can also be included in the formulation.

30 Administration of FLINT by a Nebulizer

FLINT protein can be administered by a nebulizer, such as jet nebulizer or an ultrasonic nebulizer. Typically, in

-16-

a jet nebulizer, a compressed air source is used to create a high-velocity air jet through an orifice. As the gas expands beyond the nozzle, a low-pressure region is created, which draws a solution of FLINT protein through a capillary tube connected to a liquid reservoir. The liquid stream from the capillary tube is sheared into unstable filaments and droplets as it exits the tube, creating the aerosol. A range of configurations, flow rates, and baffle types can be employed to achieve the desired performance characteristics from a given jet nebulizer. In an ultrasonic nebulizer, high-frequency electrical energy is used to create vibrational, mechanical energy, typically employing a piezoelectric transducer. This energy is transmitted to the formulation of FLINT protein either directly or through a coupling fluid, creating an aerosol including the FLINT protein. Advantageously, particles of FLINT protein delivered by a nebulizer have a particle size less than about 10 μm , preferably in the range of about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm .

Formulations of FLINT protein suitable for use with a nebulizer, either jet or ultrasonic, typically include FLINT protein in an aqueous solution at a concentration of about 1 mg to about 20 mg of FLINT protein per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and, preferably, zinc. The formulation can also include an excipient or agent for stabilization of the FLINT protein, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in such formulations include albumin, protamine, or the like. Typical carbohydrates useful in such formulations include sucrose, mannitol, lactose, trehalose, glucose, or the like. The

-17-

FLINT protein formulation can also include a surfactant, to reduce or prevent surface-induced aggregation of the FLINT protein caused by atomization of the solution in forming an aerosol. Various conventional surfactants can be employed, 5 such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbital fatty acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Additional agents known in the art for formulation of a protein can also be included in the 10 formulation.

Administration of FLINT by a Metered Dose Inhaler

In a metered dose inhaler (MDI), a propellant, FLINT protein, and any excipients or other additives are contained 15 in a canister as a mixture including a liquefied compressed gas. Actuation of the metering valve releases the mixture as an aerosol, preferably containing particles in the size range of less than about 10 μm , preferably about 1 μm to about 5 μm , and most preferably about 2 μm to about 3 μm . 20 The desired aerosol particle size can be obtained by employing a formulation of FLINT protein produced by various methods known to those of skill in the art, including jet-milling, spray drying, critical point condensation, or the like. Preferred metered dose inhalers include those 25 manufactured by 3M or Glaxo and employing a hydrofluorocarbon propellant.

Formulations of FLINT protein for use with a metered-dose inhaler device will generally include a finely divided powder containing FLINT protein as a suspension in a non 30 aqueous medium, for example, suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as

-18-

chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol and 1,1,1,2-tetrafluoroethane, HFA-134a (hydrofluroalkane-134a), HFA-227 (hydrofluroalkane-227), or the like. Preferably the propellant is a hydrofluorocarbon. The surfactant can be chosen to stabilize the FLINT protein as a suspension in the propellant, to protect the active agent against chemical degradation, and the like. Suitable surfactants include sorbitan trioleate, soya lecithin, oleic acid, or the like. In some cases solution aerosols are preferred using solvents such as ethanol. Additional agents known in the art for formulation of a protein such as FLINT can also be included in the formulation.

One of ordinary skill in the art will recognize that the methods of the current invention may be achieved by pulmonary administration of FLINT via devices not described herein.

Pharmaceutical Formulations of FLINT

The present invention also relates to a pharmaceutical composition or formulation including FLINT protein suitable for administration by inhalation. The invention also relates to methods for manufacturing formulations including FLINT protein in a form that is suitable for administration by inhalation. For example, a dry powder formulation can be manufactured in several ways, using conventional techniques. Particles in the size range appropriate for maximal deposition in the lower respiratory tract can be made by micronizing, milling, spray drying, or the like. A liquid formulation can be manufactured by dissolving the FLINT

-19-

protein in a suitable solvent, such as water, at an appropriate pH, including buffers or other excipients.

Process for Preparing Compositions of the Invention

5 Another aspect of this invention is a process for preparing a respirable powdered pharmaceutical composition of FLINT. The process comprises preparing a respirable powdered pharmaceutical composition by (a) forming a homogeneous aqueous composition comprising water, a
10 pharmaceutically acceptable excipient and FLINT suitable for treating a disease state by inhalation, (b) removing the water from the aqueous composition to form a solid and (c) transforming the resulting solid into a respirable powdered pharmaceutical composition. In another
15 embodiment, the process further comprises adding a water-soluble, physiologically-acceptable protein, such as human serum albumin (HSA), to the aqueous composition in an amount sufficient to enhance the dispersibility of the resulting respirable powdered FLINT pharmaceutical composition, as
20 described in US Patent 6,187,344 B1, herein incorporated by reference.

 In this aspect of the preparation of an aqueous mixture, a solution or stable suspension is formed by dissolving or suspending a suitable excipient, FLINT, and a
25 physiologically acceptable, water-soluble protein such as HSA in water. The order in which the components are added is not of major significance, and while the homogenous mixture may be a solution or suspension, it is preferably a solution.

30 The proportion of the components in the aqueous mixture is consistent with the proportions that are desired in the

-20-

resulting powdered composition. In general, the concentration of the materials is given in the table below:

TABLE I

5 Suitable Aqueous Compositions

	<u>Range</u>	<u>Preferred range</u>
	mg/100 ml	mg/100 ml
Excipient	15-700	500-700
FLINT	15-700	15-200
10 Water-soluble Protein	7.5-110	20-40

Another, more specific, aspect of this invention is a method for preparing a spray-dried, dispersible powdered pharmaceutical composition that comprises spray drying a homogeneous aqueous mixture comprising water, a pharmaceutically acceptable excipient, FLINT, and a dispersibility-enhancing amount of a physiologically acceptable, water-soluble protein under conditions sufficient to provide a dispersible powdered pharmaceutical composition having a particle size less than about ten microns.

Therapeutic Use for FLINT

25 The invention contemplates the administration of therapeutically effective amounts of FLINT via the pulmonary route. What constitutes a therapeutically effective amount of FLINT will depend on the particular disease or condition being treated, and the general medical condition of a patient receiving such treatment. The precise treatment dosage and regimen will depend on a variety of factors which the knowledgeable practitioner will take into account in

-21-

arriving at a particular regimen. The amount of FLINT administered may be sufficient to prevent or abate Fas-FasL induced apoptosis, which underlies many of the diseases for which FLINT is useful in treating.

5 The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

10 Generally the amount of active agent present in the composition will vary between about 0.1 % w to about 50% w., preferably from about 0.1% w. to about 5% w. and most preferably from about 0.1% w. to about 2% weight.

15 The dispersing agent useful in the composition, method and process of this invention and that provides improved dispersibility is a pharmaceutically-acceptable, water-soluble, polypeptide. For purposes of this application, polypeptide is meant to encompass both naturally occurring proteins and artificially constructed polypeptides in which
20 individual amino acid units are linked together through the standard peptide amide bond (the carboxyl group of one and the amino group of another). The dispersing agent is one that can be taken into the lungs of a patient in need thereof but will have no adverse toxicological effects at
25 the levels used. While it is preferable that the dispersing agent be an inactive agent, it is part of this invention to include agents that may have some inherent activity of their own as long as such activity is not antithetical to the utility of the overall composition. The
30 dispersing agent is characterized in having a molecular weight between about 1,000 and about 200,000. An example of an agent having a low molecular weight is a polyalanine

-22-

having a molecular weight of about 1000. Other polypeptides in that molecular weight range which are physiologically acceptable but inactive can also be prepared. Molecules that have a molecular weight in the range of about 3000 to 6000 are also useful. If a material is used that has inherent activity of its own, it is used at levels such that the inherent activity does not interfere adversely with the activity of the active agent. Another example representative of the proteins useful in this invention include alpha-lactalbumin, a constituent of milk having a molecular weight of about 14,200. Another example of a representative dispersing agent is trypsinogen, which has a molecular weight of about 24,000. A dispersing agent that is particularly preferred is human serum albumin (HSA), which has a molecular weight of about 68,000. Preferably the molecular weight of the protein dispersing agent is from about 1000 to about 100,000, particularly from about 1,000 to about 70,000 and more particularly about 68,000, i.e., HSA.

The amount of dispersing agent present in the composition of this invention may vary from about 1% w. to about 15% w., preferably from about 3% w. to about 8% w and more preferably from about 3% w to about 5% w.

In addition to the excipient of a carbohydrate, amino acid or mixtures thereof, the active agent and the protein dispersing agent, the composition of this invention may contain other pharmaceutically-acceptable excipients that may be used to stabilize the composition or make it more compatible with the unit dosage form from which it is delivered. Such excipients include, for example, buffers such as citrate, phosphate or acetate.

-23-

The composition of this invention will be delivered from a unit dosage receptacle containing an amount that will be sufficient to provide the desired physiological effect upon inhalation by a subject in need thereof. The amount will be dispersed in a chamber that has an internal volume sufficient to capture substantially all of the powder dispersion resulting from the unit dosage receptacle. Usually the volume of the chamber will be from about 50 ml to about 1000 ml, preferably from about 100 ml to about 750 ml. Thus, the unit dosage amount will be from about 2 mg of powder to about 20 mg of powder preferably about 4 mg to about 10 mg of powder per unit dosage. About 5 mg per unit dosage is quite effective. The preferred unit dosage receptacle is a blister pack, generally provided as a series of blister pack strips. The general process for preparing such blister packs or blister pack strips is generally known to one of skill in the art from such publications as Remington's Pharmaceutical Sciences (18th Edition) or other similar publications. The volume of such dosage form receptacle to accommodate the needed amount of powder of this invention will be about 1 ml to about 30 ml, preferably about 2 ml to about 10 ml.

25

Example 1

Serum Pharmacokinetics of Human FLINT in Beagle Dogs
Following Pulmonary Administration of Single Aerosolized
Doses

Aerosols of human FLINT generated from solutions of FLINT in sterile water, are administered to anesthetized dogs by the pulmonary route through an endotracheal tube via an ultrasonic nebulizer. Serum concentration of

30

-24-

immunoreactive FLINT are determined by validated radioimmunoassay methods.

Six beagle dogs (3 male and 3 female) are used in the study. Animals are housed either two per cage or
5 individually in stainless steel cages with suspended mesh floors. Initially, all dogs are fed approximately 450 g of Purina Certified Canine Diet 5007 each day. Animals are fasted approximately eight hours before dosing. After recovery from anesthesia, food and water are provided *ad*
10 *libitum* until 48 hours post-dose. The initial daily feeding regimen is initiated at 48 hours post-dose.

Blood samples are collected at various time points after dosing to determine plasma concentrations of the human FLINT and bioavailability of inhaled material is determined.
15 Dogs are chosen because they are large animals with respiratory tract deposition of particles similar to man.

The effectiveness of pulmonary administration of FLINT is indicated by increased concentrations of immunoreactive FLINT in the serum of treated dogs.
20

Example 2

Serum Pharmacokinetics of FLINT Analog R218Q in Beagle Dogs

Following Pulmonary Administration of Single Aerosolized

25 Doses

Aerosols of human FLINT analog R218Q are generated from a solution of R218Q in sterile water, and are administered to anesthetized dogs by the pulmonary route through an endotracheal tube via an ultrasonic nebulizer. Serum
30 concentration of immunoreactive FLINT R218Q are determined by validated radioimmunoassay methods.

-25-

Six beagle dogs (3 male and 3 female) are used in the study. Animals are housed either two per cage or individually in stainless steel cages with suspended mesh floors. Initially, all dogs are fed approximately 450 g of
5 Purina Certified Canine Diet 5007 each day. Animals are fasted approximately eight hours before dosing. After recovery from anesthesia, food and water are provided *ad libitum* until 48 hours post-dose. The initial daily feeding regimen is initiated at 48 hours post-dose.

10 Blood samples are collected at various time points after dosing to determine plasma concentrations of the FLINT R218Q and bioavailability of inhaled material is determined.

The effectiveness of pulmonary administration of FLINT R218Q is indicated by increased concentrations of
15 immunoreactive FLINT R218Q in the serum of treated dogs.

-26-

What is claimed is:

- 5 1. A method comprising pulmonary administration of a therapeutically effective amount of FLINT or FLINT analog.
2. A method as in claim 1 wherein said FLINT is delivered to
10 a lower airway of a patient.
3. A method as in claim 2 wherein said FLINT is deposited in the alveoli.
- 15 4. A method as in claim 1 wherein said FLINT is inhaled through the mouth of a patient.
5. A method as in claim 1, said FLINT further comprising a pharmaceutically acceptable carrier to form a FLINT
20 composition.
6. A method as in claim 5 wherein said composition is selected from the group consisting of an aqueous solution, an aqueous suspension, a non-aqueous
25 suspension, or a dry powder.
7. A method of claim 6 wherein said composition is administered as an aerosol.
- 30 8. A method of claim 6 wherein said composition is a dry powder.

-27-

9. A method as in claim 5 wherein said FLINT has a particle size of less than about 10 microns.
10. A method as in claim 5 wherein said FLINT has a particle size of about 1 micron to about 5 microns.
11. A method as in claim 5 wherein said FLINT has a particle size of about 2 microns to about 3 microns.
- 10 12. A method as in claim 1 wherein said FLINT is delivered from an inhalation device suitable form pulmonary administration and capable of depositing FLINT in the lungs of a patient.
- 15 13. A method of claim 12 wherein said device is selected from the group consisting of a nebulizer, a metered-dose inhaler, a dry-powder inhaler and a sprayer.
14. A method of Claim 1 wherein said analog is R218Q.

SEQUENCE LISTING

<110> Eli Lilly and Company

<120> PULMONARY ADMINISTRATION OF FLINT

<130> X-13678

<160> 5

<170> PatentIn version 3.0

<210> 1

<211> 271

<212> PRT

<213> Homo sapiens

<400> 1

```

Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu
1          5          10          15

Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro
20          25          30

Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His
35          40          45

Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val
50          55          60

Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His
65          70          75          80

Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe
85          90          95

Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro
100         105         110

Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr
115         120         125

Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn
130         135         140

Cys Thr Ala Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His
145         150         155         160

Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val
165         170         175

Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe
180         185         190

Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu
195         200         205

Ala Pro Glu Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu
210         215         220

```

Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp
 225 230 235 240

Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met
 245 250 255

Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His
 260 265 270

<210> 2
 <211> 813
 <212> DNA
 <213> Homo sapiens

<400> 2
 gtggcagaaa caccaccta ccctggcgg gacgcagaga caggggagcg gctggtgtgc
 60

gcccagtgcc cccaggcac ctttgtgcag cggccgtgcc gccgagacag cccacgcacg
 120

tgtggcccgt gtccaccgcg cactacacg cagttctgga actacctgga gcgctgccgc
 180

tactgcaacg tcctctgcgg ggagcgtgag gaggaggcac gggcttgcca cgccaccac
 240

aaccgtgcct gccgctgcg caccggcttc ttgcgcacg ctggtttctg cttggagcac
 300

gcatcgtgtc cacctggtgc cggcgtgatt gccccgggca cccccagcca gaacacgcag
 360

tgcagccgt gccccccagg caccttctca gccagcagct ccagctcaga gcagtgccag
 420

ccccaccgca actgcacggc cctgggcctg gccctcaatg tgccaggctc ttcctccat
 480

gacaccctgt gcaccagctg cactggcttc ccctcagca ccagggtacc aggagctgag
 540

gagtgtgagc gtgccgtcat cgactttgtg gctttccagg acatctccat caagaggctg
 600

cagcggctgc tgcaggccct cgaggccccg gagggctggg gtccgacacc aagggcgggc
 660

cgcgccgcct tgcagctgaa gctgcgtcgg cggctcacgg agctcctggg ggcgcaggac
 720

ggggcgctgc tgggtcggct gctgcaggcg ctgcgcgtgg ccaggatgcc cgggctggag
 780

cggagcgtcc gtgagcgctt cctccctgtg cac
 813

<210> 3

<211> 300
 <212> PRT
 <213> Homo sapiens

<400> 3

```

Met Arg Ala Leu Glu Gly Pro Gly Leu Ser Leu Leu Cys Leu Val Leu
1          5          10          15

Ala Leu Pro Ala Leu Leu Pro Val Pro Ala Val Arg Gly Val Ala Glu
          20          25          30

Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu Arg Leu Val
          35          40          45

Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro Cys Arg Arg
          50          55          60

Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His Tyr Thr Gln
65          70          75          80

Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val Leu Cys Gly
          85          90          95

Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His Asn Arg Ala
          100          105          110

Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe Cys Leu Glu
          115          120          125

His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro
          130          135          140

Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala
145          150          155          160

Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala
          165          170          175

Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His Asp Thr Leu
          180          185          190

Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala
          195          200          205

Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile
210          215          220

Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu
225          230          235          240

Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys
          245          250          255

Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu
          260          265          270

Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu
          275          280          285

```


Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His
 290 295 300

<210> 4
 <211> 936
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (25)..(924)

<400> 4
 gctctccctg ctccagcaag gacc atg agg gcg ctg gag ggg cca ggc ctg
 51
 Met Arg Ala Leu Glu Gly Pro Gly Leu
 1 5

tcg ctg ctg tgc ctg gtg ttg gcg ctg cct gcc ctg ctg ccg gtg ccg
 99
 Ser Leu Leu Cys Leu Val Leu Ala Leu Pro Ala Leu Leu Pro Val Pro
 10 15 20 25

gct gta cgc gga gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca
 147
 Ala Val Arg Gly Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala
 30 35 40

gag aca ggg gag cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt
 195
 Glu Thr Gly Glu Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe
 45 50 55

gtg cag cgg ccg tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt
 243
 Val Gln Arg Pro Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys
 60 65 70

cca ccg cgc cac tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc
 291
 Pro Pro Arg His Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg
 75 80 85

tac tgc aac gtc ctc tgc ggg gag cgt gag gag gag gca cgg gct tgc
 339
 Tyr Cys Asn Val Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys
 90 95 100 105

cac gcc acc cac aac cgt gcc tgc cgc tgc cgc acc ggc ttc ttc gcg
 387
 His Ala Thr His Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala
 110 115 120

cac gct ggt ttc tgc ttg gag cac gca tcg tgt cca cct ggt gcc ggc
 435
 His Ala Gly Phe Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly
 125 130 135

gtg att gcc ccg ggc acc ccc agc cag aac acg cag tgc cag ccg tgc
 483
 Val Ile Ala Pro Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys
 140 145 150

ccc cca ggc acc ttc tca gcc agc agc tcc agc tca gag cag tgc cag
 531
 Pro Pro Gly Thr Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln
 155 160 165

ccc cac cgc aac tgc acg gcc ctg ggc ctg gcc ctc att gtg cca ggc
 579
 Pro His Arg Asn Cys Thr Ala Leu Gly Leu Ala Leu Ile Val Pro Gly
 170 175 180 185

tct tcc tcc cat gac acc ctg tgc acc agc tgc act ggc ttc ccc ctc
 627
 Ser Ser Ser His Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu
 190 195 200

agc acc agg gta cca gga gct gag gag tgt gag cgt gcc gtc atc gac
 675
 Ser Thr Arg Val Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp
 205 210 215

ttt gtg gct ttc cag gac atc tcc atc aag agg ctg cag cgg ctg ctg
 723
 Phe Val Ala Phe Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu
 220 225 230

cag gcc ctc gag gcc ccg gag ggc tgg gct ccg aca cca agg gcg ggc
 771

Gln Ala Leu Glu Ala Pro Glu Gly Trp Ala Pro Thr Pro Arg Ala Gly

235

240

245

cgc gcg gcc ttg cag ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg
819

Arg Ala Ala Leu Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu

250

255

260

265

ggg gcg cag gac ggg gcg ctg ctg gtg cgg ctg ctg cag gcg ctg cgc
867

Gly Ala Gln Asp Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg

270

275

280

gtg gcc agg atg ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc
915

Val Ala Arg Met Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu

285

290

295

cct gtg cac tgatcctggc cc
936

Pro Val His

300

<210> 5

<211> 300

<212> PRT

<213> Homo sapiens

<400> 5

Met Arg Ala Leu Glu Gly Pro Gly Leu Ser Leu Leu Cys Leu Val Leu
1 5 10 15

Ala Leu Pro Ala Leu Leu Pro Val Pro Ala Val Arg Gly Val Ala Glu
20 25 30

Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu Arg Leu Val
35 40 45

Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro Cys Arg Arg
50 55 60

Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His Tyr Thr Gln
65 70 75 80

Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val Leu Cys Gly
 85 90 95

Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His Asn Arg Ala
 100 105 110

Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe Cys Leu Glu
 115 120 125

His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro
 130 135 140

Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala
 145 150 155 160

Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala
 165 170 175

Leu Gly Leu Ala Leu Ile Val Pro Gly Ser Ser Ser His Asp Thr Leu
 180 185 190

Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala
 195 200 205

Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile
 210 215 220

Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu
 225 230 235 240

Gly Trp Ala Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys
 245 250 255

Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu
 260 265 270

Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu
 275 280 285

Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His
 290 295 300

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/09668 A3

(51) International Patent Classification⁷: **A61K 9/00**

(21) International Application Number: **PCT/US01/21105**

(22) International Filing Date: **20 July 2001 (20.07.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/222,476 **2 August 2000 (02.08.2000)** **US**

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY [US/US]**; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MICANOVIC, Radmila [US/US]**; 7126 White Oak Trail, Indianapolis, IN 46236 (US). **WITCHER, Derrick, Ryan [US/US]**; 10898 Parrot Court, Fishers, IN 46038 (US).

(74) Agents: **WEBSTER, Thomas, D. et al.**; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).

(81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**

(84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
10 May 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/09668 A3

(54) Title: **PULMONARY ADMINISTRATION OF FLINT**

(57) Abstract: The claimed invention relates to methods and compositions for administering FLINT by inhalation, and a methods for treating a disease by administering FLINT by inhalation.

INTERNATIONAL SEARCH REPORT

Internat. Application No PCT/US 01/21105		
A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K9/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) WPI Data, PAJ, EPO-Internal, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 37094 A (ELI LILLY) 29 June 2000 (2000-06-29) claims 1,6,10 page 42, line 5 -page 43, line 5 page 44, line 21 - line 30 -----	1,4-7, 12,13
<div style="display: flex; justify-content: space-between;"> <div> <input type="checkbox"/> Further documents are listed in the continuation of box C. </div> <div> <input checked="" type="checkbox"/> Patent family members are listed in annex. </div> </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents:</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">19 March 2002</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">26/03/2002</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Ventura Amat, A</div>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21105

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0037094	A	29-06-2000	
		AU 2211100 A	12-07-2000
		AU 3369199 A	18-10-1999
		BR 9909328 A	12-12-2000
		CA 2324517 A1	07-10-1999
		CN 1303429 T	11-07-2001
		CZ 20003433 A3	17-10-2001
		EP 1140138 A2	10-10-2001
		HU 0102067 A2	28-10-2001
		NO 20004873 A	24-11-2000
		PL 343847 A1	10-09-2001
		TR 200002824 T2	21-12-2000
		WO 9950413 A2	07-10-1999
		WO 0037094 A2	29-06-2000